

Coral reefs are “holobionts,” composed of coral-forming species, numerous algae, and bacterial species that live symbiotically with their coral hosts (Roik et al., 2022). Predominantly found in tropical and subtropical waters of the Atlantic, Pacific, and Indian Oceans, these holobionts serve vital roles in marine ecosystems, acting as protective barriers and food reservoirs for approximately one third of marine species (Plaisance, 2011). Among those marine species are coral-dwelling fish, which one-third gain 80% of their diet from coral reefs (Cole et al., 2008). Other fish species utilize coral reefs as breeding grounds and nurseries for their young (Verweij et al., 2008). Despite overwhelming evidence suggesting fish utilize coral reefs for multifold purposes, it is unknown the extent to which they would be impacted if coral reefs went extinct.

Some coral species act as ecosystem engineers, such as those belonging to the genera Scleractinia, by secreting calcium carbonate to build coral reef structures (Von Euw et al., 2017). However, this process may be halted due to environmental change. Low oceanic pH or high ocean temperatures cause available oceanic carbonate ions to be converted to carbonic acid (Mollica et al., 2018). With less carbonate ions available, corals

6RPH RIWKH YW FRDQ QDPHQW BPDWRSRBLGH D H WKRJKW
WR KDYH HYRQHG GKKJ WKH ZGRYLF LDQ 3HURG QHDU PLORQH DW
ago (Webby, 2002). During this time, corals transitioned from microbial-based metazoan symbionts to algal and microbial-based symbionts (Webby, 2002). While the Ordovician is surely an important marking for the first appearance of coral species, the Devonian Period displays mass-scale population shifts which are of interest to coral extinction literature. During the Devonian, coral species faced mass expansion, the formation of the first coral reef networks, and the demise of such networks (Wendt & Kaufmann, 2006, Bridge et al., 2022). Overall, the his-

explosion of aquatic life, thus commonly referred to as “The Age of Fishes”. Alongside other marine life, coral life multiplied in increasing numbers, with one of the largest reef complexes being created by the middle of the Devonian Period (Wendt & Kaufmann, 2006). By the end of the period, these same reef ecosystems collapsed, a period referred to as the Late Devonian Mass Extinction (Bridge et al., 2022). The exact factors that may have led to this collapse are unknown, although eutrophication, anoxia, hypersalinity, red tide and algal blooming may have contributed (Gong et al., 2002). Nonetheless, while the Devonian holds remarkable developments in coral-reef complexes, it also holds the extinction of coral reef species in such complexes.

Coral reefs have faced considerable threats between the Devonian Period and the Quaternary Period of the Anthropocene. Facing comet showers, volcanic eruptions, and glacial and interglacial periods during the Devonian, corals predominantly face disease and anthropogenic effects today. The similar threat to corals of global warming transverses through both periods, but it is accelerated today due to anthropogenic effects like the burning of fossil fuels. The effects of global warming on coral reefs have been studied for decades, yet literature discussing the implications of coral reef's mortality on marine ecosystems remains underserved.

Coral reefs house nearly 600,000 to 9 million species worldwide (Plaisance, 2011). From those, coral-dwelling fish species rely extensively on coral-reefs for protection from predators and access to food (Plaisance, 2011, Cole et al., 2008). Therefore, it is important to project how marine communities, specifically coral-dwelling fish species, might respond if coral extinction were to occur in the future. Some fish species are obligate coral species, meaning they rely on corals for survival, whereas other species earn facultative benefits from corals. Therefore, the consequences of coral extinction might differ among marine ecosystems.

Fish primarily utilize coral reefs by using their trabeculated coral branches as shelters from larger predators (Boström-Einarsson et al., 2018). However, as coral branches die, they become overgrown with algae, thereby minimizing available space for hiding and shelter (Boström-Einarsson et al., 2018). Researchers found as available shelter decreases, fish are more likely to seek live coral, crossing large areas of open water between coastlines and increasing their likelihood of encountering a predator (Boström-Einarsson et al., 2018). They found a positive relationship between coral death and predation-induced mortality of coral-reef dwelling damselfish, *Pomacentrus moluccensis* (Boström-Einarsson et al., 2018). The loss of available trabeculated coral branches may be particularly harmful for the 126 species of coral-dwelling fish who use coral branches as shelter from predators during their infancy (Coker et al., 2013, Jones et al., 2004). In the case of coral extinction, we may use these findings to suggest coral-reef dwelling fish populations would suffer higher mortality rates due to their increased risk of finding shelter in open waters.

It is also suggestive that coral reef fish communities would be reshaped completely if coral reefs went extinct, potentially threatening species abundance and ecosystem functioning (Jones et al., 2004, Cheal et al., 2008). Fish species diversity at the Great Barrier Reef was measured over an eleven-year period (Cheal et al., 2008). Despite coral abundance decreasing following coral bleaching and mortality events, diversity remained the same (Cheal et al., 2008). It is important to note, however, community diversity is not necessarily indicative of communal change in species presence. As coral mortality increased, so too did the abundance of herbivorous fish who do not depend on corals for survival (Cheal et al., 2008). Coral-dependent fish species and other fish species abundance declined with coral mortality, thereby reducing overall ecosystem functioning (Cheal et al., 2008). This shows that coral bleaching and coral decline can reshape fish species abundance at coral reefs, reducing the extent to which coral reefs can provide natural services and products to the surrounding environment (Cheal et al., 2008). Overall, although community diversity (the number of coral-dwelling fish interacting at coral reefs) may not decline, species comv39vi0 0 8 36 293.6g at PAVLHFWV0FRDGZHCQYK0QRW0

Areas Improve Reef Resilience? *Annual Review of Marine Science*, 11(1), 307–334.

Buddemeier, R. W., & Fautin, D. G. 1993. Coral bleaching as an Adaptive Mechanism. *BioScience*, 43(5), 320–326.

Cerrano, C., Cardini, U., Bianchelli, S., Corinaldesi, C., Pusceddu, A., & Danovaro, R. 2013. Red Coral extinction risk enhanced by ocean acidification. *Scientific Reports*, 3(1).

Cheal, A. J., Wilson, S. K., Emslie, M. J., Dolman, A. M., & Sweatman, H. 2008. Responses of reef fish communities to coral declines on the Great Barrier Reef. *Marine Ecology Progress Series*, 372, 211–22

Chen, D., Qing, H., & Li, R. 2005. The late devonian frasnian–famennian (f/F) biotic crisis: Insights from C, C and Sr/SR Isotopic Systematics. *Earth and Planetary Science Letters*, 235(1-2), 151–166.

Coker, D. J., Wilson, S. K., & Pratchett, M. S. 2013. Importance of live coral habitat for reef fishes. *Reviews in Fish Biology and Fisheries*, 24(1), 89–126.

Cole, A. J., Pratchett, M. S., & Jones, K. P. 2008. The impact of coral reef degradation on fish diversity and abundance. *Marine Ecology Progress Series*, 372, 211–22