

Oxidation rates of apple slices under

different pH solutions

Holly Kiernan

hkiernan@lafc.edu

hkiernan@lafc.edu

Introduction:

When an apple is cut, the enzyme polyphenol oxidase (PPO) reacts with oxygen, now present in plant cells, to produce brown pigments called polyphenols.

The rate of browning is affected by the pH of the solution. In an acidic solution (pH 2), the PPO enzyme is denatured and browning is slowed. In a basic solution (pH 10), the PPO enzyme is more active and browning is accelerated.

The purpose of this experiment is to determine the effect of pH on the rate of apple browning. The experiment will be conducted using three solutions: pH 2 (lemon juice), pH 7 (distilled water), and pH 10 (baking soda solution).

It is hypothesized that the rate of browning will be lowest in the pH 2 solution and highest in the pH 10 solution. The pH 7 solution will serve as a control.

Hypothesis and Prediction:

The hypothesis is that pH will have an impact on apple oxidation. The more acidic the solution that the apple is placed into, the less browning will be observed due to the pH of the solution leading to the denaturing of the enzyme PPO. Lemon juice, an acid with a pH of about 2, is commonly used to prevent apple oxidation, providing support for the hypothesis that an acidic solution will slow browning more than a basic solution. The solution of pH 2 acted as the acidic solution that should reduce the rate of apple oxidation. Bases like baking soda, for example, are not typical used to slow apple browning like lemon juice is. Suppose apple slices are placed in an acidic solution (pH 2) will exhibit a slower rate of oxidation over the course of 180 minutes by appearing less brown. The solution of pH 2 will denature the enzymes to the greatest extent, while pH 7 will denature the enzymes to a lesser extent, and pH 10 will not denature the enzymes. Finally, the browning rates of apples exposed to air will be higher than all the other apples placed in solutions because oxygen will be able to enter the damaged apple cells more easily. However, apples placed in water will display similar oxidation rate to the apples in pH 10. dH2O is neutral, but the water will help keep oxygen from entering the apple cells.

Materials and Methods:

The experiment was conducted using three solutions: pH 2 (lemon juice), pH 7 (distilled water), and pH 10 (baking soda solution). Apples were cut into slices and placed in each solution. The color of the apple slices was measured at 0, 30, 60, 90, 120, 150, and 180 minutes.

For the four cups that contained the solution, enough of the solution was added to cover the apples. For each condition, about 5mL of solution was enough to cover the apples. In order to observe oxidation, the apples were kept in the cups and allowed to soak in their respective solutions. The color of the apple slices was measured at 0, 30, 60, 90, 120, 150, and 180 minutes. The color of the apple was matched to the closest corresponding value, and the average color of the condition at every time point was calculated to examine the oxidation rate of apples (Figure 1).



Figure 1. The color of the apple slice correlates to a deeper brown color and, thus, a higher rate of melanin production and apple oxidation. In other words, the darker the color, the more the apple slice has oxidized. This color palette, thus, was used in order to quantify apple browning data.

Written Results:

Initially, the average color value for all the apple slices across all conditions was 1.3. The pH 2 solution, however, stayed at a color value of 1, with no browning being observed. At the 30-minute mark, the pH 7 solution had a color value of 1.3, and the pH 10 solution had a color value of 3.3. The pH 2 solution, however, stayed at a color value of 1.3. The dH2O solution had a color value of 1.3. The pH 2 solution, however, stayed at a color value of 1.3. The pH 7 solution had a color value of 1.3, and the pH 10 solution had a color value of 3.3. The pH 2 solution, however, stayed at a color value of 1.3. The pH 7 solution had a color value of 1.3, and the pH 10 solution had a color value of 3.3. The pH 2 solution, however, stayed at a color value of 1.3. The pH 7 solution had a color value of 1.3, and the pH 10 solution had a color value of 3.3.

solutions like lemon juice have been observed to slow melanin production in apples, and so that knowledge went into forming the hypothesis. As evidenced, PPO activity was not impacted, at least all the apple slices in the solution, if not all apple slices, would all have shown the same average color value. The data, however, shows a wide range of color values, with a higher color value indicating higher PPO activity and, thus, greater melanin production. Solutions like lemon juice have been observed to slow melanin production in apples, and so that knowledge went into forming the hypothesis. As evidenced, PPO activity was not impacted, at least all the apple slices in the solution, if not all apple slices, would all have shown the same average color value. The data, however, shows a wide range of color values, with a higher color value indicating higher PPO activity and, thus, greater melanin production.

Figure 2. A line graph showing the average color value of apple slices over time (0 to 30 minutes) for three different pH solutions: pH 2 (acidic), pH 7 (neutral), and pH 10 (basic). The y-axis represents the average color value (10 to 50), and the x-axis represents time in minutes (0 to 30). The pH 2 solution shows the highest color value, followed by pH 7 and then pH 10. Error bars represent standard deviation.

Time (min)	pH 2 (Acidic)	pH 7 (Neutral)	pH 10 (Basic)
0	10	10	10
5	15	12	10
10	20	15	12
15	25	18	15
20	30	20	18
25	35	22	20
30	40	25	22

Conclusion:

The data found in this experiment supported the hypothesis that pH of the solution the apple is in. It was previously predicted that a lower pH would lead to less melanin production or a slower oxidation rate because the acid would denature the enzyme polyphenol oxidase. The prediction was partly correct, with the apples in a pH 2 solution having a higher average color value than the solution of pH 10. However, the scoring on the apples immersed in pH 10 solution also showed melanin production because oxygen could not react with PPO. Only acidic solutions may be able to slow the rate of oxidation in comparison to mildly acidic solutions possibly explaining the average color values being somewhat close to each other. Thus, the hypothesis is still supported due to similar melanin production in neutral conditions in both cases. However, in the case of a basic solution, the oxygen from entering some of the damaged apple cells. The apples in the air have no protection from oxygen, in other words.

The pH 2 and pH 10 solutions were chosen to have a spectrum of acidic to neutral to basic solutions. As mentioned before, acidic

solutions like lemon juice have been observed to slow melanin production in apples, and so that knowledge went into forming the hypothesis. As evidenced, PPO activity was not impacted, at least all the apple slices in the solution, if not all apple slices, would all have shown the same average color value. The data, however, shows a wide range of color values, with a higher color value indicating higher PPO activity and, thus, greater melanin production.