Dengue Fever in China

Jordan Garland
Lake Forest College
Lake Forest, Illinois 60045

with 13 deaths occurring in the same 15-year period (Yue et al., 2022).

Additionally, there are some groups that are more likely to fall ill with dengue fever globally. For instance, those living in low-income communities lacking medical supplies, treatment, and intervention are more likely to get sick and die from dengue. Also, the most severe cases were reported in children under 15 years of age throughout hyperendemic areas in Asia, including China (Tantawichien, 2012). There is currently no specifc antiviral for dengue, but the symptoms can be managed, sub

Control and Prevention, 2019). This vaccine is efective, with an efcacy of approximately 80 percent. It is, however, only administered to individuals from 12 to 5] , ˈ oni ſ Ə ˈ ini indi

risk of giving someone severe dengue if they did not have it previously.

In China specifcally, research has proven that socioeconomic factors (defned as interactions between social and economic factors) signifcantly afect how damaging dengue can be. For example, population density, urbanization, and gross domestic product (GDP) can alter the

are known to impact dengue incidence. In spite of this, it is shown that there are some interventions that are proven to be efective, including

communities are particularly at risk. As such, funding is necessary to study dengue further and implement efective preventative interventions.

Throughout Chinr r s ɔ̈ l ą ɹ m ƥ ƫ T m Štd ʊ m ʊ E T fuu u m MstueMs , ƫ

examined. The data concluded that a higher population density comes with a higher conditional probability of the risk of disease incidence. This higher risk can result from the fact that with more people in each area, there is a higher likelihood for infected mosquitos to bite people. On the other hand, with GDP per capita, the evidence showed the opposite. With less GDP, a region is at higher risk for dengue incidence. Since a lower GDP is associated with lower income, corresponding areas are less likely to have interventions in place or the required resources for treatment, increasing risk. Higher NDVI, which quantifes the presence of vegetation in a given area, was shown to decrease disease incidence. This conclusion seemingly goes against preconceived notions. One would think that more forests and vegetation correspond with more mosquitoes and, therefore, incidence. This was not the case, however. Finally, the data on higher road density shows an increase in the risk of disease incidence. An area with more roads in

In further support of these ideas, another research article studied the efects of natural and socioeconomic factors on dengue fever in two cities in China, intending to identify the key factors afecting dengue fever's spatial and temporal distribution (Chen et al., 2020). The two provinces were Foshan and Guangzhou, areas known to have had severe dengue outbreaks in

relative risk was calculated. Experimenters heavily scrutinize population density. In this 12-year period, Chen and company examine the relative risk

sity, those being central Guangzhou, Eastern Foshan, and the borders between the two areas, are shown to have a higher risk for dengue incidence.

tion and density correlate with a higher relative risk for dengue disease and incidence. The most vu / signifcance that both extremely high temperatures a

tiveand its evolution over the past 30 years by compiling data from the Global

Burden of Diseases, Injuries and Risk Factors Study 2019 (GBD 2019). This study examines four areas of interest: the age-standardized incidence rate (ASIR) or disease incidence, the age-standardized death rate (ASDR) or mortality, and age-standardized disability—adjusted life years. (DALYs) The last of these is the estimated number of years of life lost from dengue. China had exceptionally high levels of ASIR and DALYs, with a value of approximately three to six (in units of 100,000) for the former and a value of four to eight years of life lost for the latter. Likewise, the same study reported that East Asia, which includes China had the fastest rise of ASIR